

- [4] Ward E, Boffetta P, Andersen A, Colin D, Comba P, Deddens JA, et al. Update of the follow-up of mortality and cancer incidence among European workers employed in the vinyl chloride industry. *Epidemiology* 2001;12:710–718.
- [5] Mastrangelo G, Fedeli U, Fadda E, Valentini F, Agnesi R, Magarotto G, et al. Increased risk of hepatocellular carcinoma and liver cirrhosis in vinyl chloride workers: synergistic effect of occupational exposure with alcohol intake. *Environ Health Perspect* 2004;112:1188–1192.
- [6] Boffetta P, Matisane L, Mundt KA, Dell LD. Meta-analysis of studies of occupational exposure to vinyl chloride in relation to cancer mortality. *Scand J Work Environ Health* 2003;29:220–229.
- [7] Pirastu R, Baccini M, Biggeri A, Comba P. Epidemiologic study of workers exposed to vinyl chloride in Porto Marghera: mortality update. *Epidemiol Prev* 2003;27:161–172.
- [8] Ryder SD. Guidelines for the diagnosis and treatment of hepatocellular carcinoma (HCC) in adults. *Gut* 2003;52 (Suppl. 3):1–8.
- [9] Dragani TA, Zocchetti C. Occupational exposure to vinyl chloride and risk of hepatocellular carcinoma. *Cancer Causes Control* 2008;19:1193–1200.
- [10] McLaughlin JK, Lipworth L. A critical review of the epidemiologic literature on health effects of occupational exposure to vinyl chloride. *J Epidemiol Biostat* 1999;4:253–275.
- [11] Grosse Y, Baan R, Straif K, Secretan B, El Ghissassi F, Bouvard V, et al. Carcinogenicity of 1,3-butadiene, ethylene oxide, vinyl chloride, vinyl fluoride, and vinyl bromide. *Lancet Oncol* 2007;8:679–680.

Giuseppe Mastrangelo*

Department of Environmental Medicine and Public Health,
University of Padova, Padova, Italy

* Corresponding author. Address: University of Padova,
Via Giustiniani 2, 35128 Padova, Italy.
Tel.: +39 049 8212543; fax: +39 049 8212542.
E-mail address: giuseppe.mastrangelo@unipd.it

Diego Martines

Department of Surgical and Gastroenterological Sciences,
University of Padova, Padova, Italy

Ugo Fedeli

SER-Epidemiological Department,
Veneto Region, Castelfranco Veneto, Italy

Selective rather than routine approach to endoscopic retrograde cholangio-pancreatography in diagnosis of biliary atresia

To the Editor:

Petersen and colleagues describe their experience in the “routine” use of endoscopic retrograde cholangio-pancreatography (ERCP) in children awaiting explorative laparotomy (EL) for suspected biliary atresia (BA) [1]. They report that almost 25% of the infants have avoided surgery after documenting a patent biliary tree. The quoted specificity of ERCP in diagnosing BA was 73% [1].

The aim of avoiding unnecessary surgery is valiant and undoubtedly, but we feel that a non-selective approach to a relatively invasive procedure in jaundiced infants, who may be recovering from the potentially recoverable liver injury, such as the one secondary to prematurity, total parenteral nutrition, severe haemolysis and/or infection, may not be justified. Repeated general anaesthetic, required by combined ERCP/EL, in a jaundiced child carries additional risks outside the usual economic arguments. Sixty (43%) infants referred to Petersen et al. who underwent ERCP actually did not have BA. Suspected BA may mean different things for different centres and submitting all referred children to ERCP may not be indicated, despite the technical excellence.

In this study the “pre-selection” information, including ultrasound findings or liver histology, was unfortunately not presented, although the authors have quoted experiences where specificity for percutaneous liver biopsy was 96% [2] and 89% [3], respectively. An early study from our centre found the liver histology as sufficient for diagnosis of BA in 86% infants with neonatal cholestasis [4]. Finally, in the study by Petersen et al. the children undergoing ERCP/EL combination were up to 174 days old, an age where benefits of corrective surgery are somewhat dubious.

We have recently reported our own tertiary centre experience on use of ERCP in 48 cholestatic infants younger than 100 days, representing only around 4% of infants with neonatal cholestasis, where diagnosis after comprehensive hepatological work-up remained unclear [5]. EL was avoided in 42% of children, while selective ERCP had a specificity of 87% for diagnosis of BA [5].

Protocols for investigating infants with prolonged neonatal cholestasis will continue to differ from centre to centre, but there is no substitute for evaluation of all the available clinical information, including monitoring stool colour and its change on choleretics [6]. ERCP is a very welcome addition to the diagnostic algorithm for the diagnosis of BA, but in our view should be used selectively once other, less invasive, tests have proven inconclusive.

References

- [1] Petersen C, Meier PN, Schneider A, Turowski C, Pfister ED, Manns MP, et al. Endoscopic retrograde cholangiopancreatography prior to explorative laparotomy avoids unnecessary surgery in patients suspected for biliary atresia. *J Hepatol* 2009;51:1055–1060.
- [2] Park WH, Choi SO, Lee HJ, Kim SP, Zeon SK, Lee SL. A new diagnostic approach to biliary atresia with emphasis on the ultrasonographic triangular cord sign: comparison of ultrasonography, hepatobiliary scintigraphy, and liver needle biopsy in the evaluation of infantile cholestasis. *J Pediatr Surg* 1997;32:1555–1559.
- [3] Esmaili J, Izadyar S, Karegar I, Gholamrezaezhad A. Biliary atresia in infants with prolonged cholestatic jaundice. diagnostic accuracy of hepatobiliary scintigraphy. *Abdom Imaging* 2007;32:243–247.
- [4] Manolaki AG, Larcher VF, Mowat AP, Barrett JJ, Portmann B, Howard ER. The prelaparotomy diagnosis of extrahepatic biliary atresia. *Arch Dis Child* 1983;58:591–594.
- [5] Shanmugam NP, Harrison PM, Devlin J, Peddu P, Knisely A, Davenport M, et al. Selective use of endoscopic retrograde cholangiopancreatography in the diagnosis of biliary atresia in infants younger than 100 days. *J Pediatr Gastroenterol Nutr* 2009;49:435–441.
- [6] Hadzic N. Biliary atresia. In: Jara P, editor. Liver diseases in children. International hepatology updates. Barcelona: Permanyer Publications; 2009.

Nedim Hadzic,

Phillip M. Harrison

King's College Hospital, London, United Kingdom

E-mail address: nedim.hadzic@kcl.ac.uk